

IMPACT: AIR POLLUTANTS EMISSIONS

Reference 13

Wang, Y; Dong, HM; Zhu, ZP; Gerber, PJ; Xin, HW; Smith, P; Opio, C; Steinfeld, H; Chadwick, D 2017 Mitigating Greenhouse Gas and Ammonia Emissions from Swine Manure Management: A System Analysis ENVIRONMENTAL SCIENCE & TECHNOLOGY 10.1021/acs.est.6b06430

Background and objective

Studies have been conducted to address manure-related emissions, and various mitigation measures have been tested and developed. However, most studies have focused either on one specific gas, one individual manure management phase or influencing factor, or mitigation practice. The objective of this study is to estimate the emissions mitigation potentials for NH₃, methane (CH₄), and nitrous oxide (N₂O) of different swine manure storage and treatment mitigation strategies. Here the results concerning NH₃ emissions are reported.

Search strategy and selection criteria

The ISI Web of Knowledge database (www.isiwebofknowledge.com) and the Chinese journal database (www.cnki.net) were used to search all published data sets as of January 2016. Specific search terms were combined and used, depending on animal categories (swine, pig, livestock, animal), manure, in-house manure management (slatted floor, pit, bedding, litter, pull-plug, discharge, scraper, separation), outdoor manure management (lagoon, slurry pond, storage tank, compost, solid storage, stockpile), land application (surface spreading, injection, incorporation, band spreading), gaseous emission (NH₃, CH₄, N₂O, and GHG gas), and mitigation measure (diet, biofilter, biogas, additive, cover, acid, cooling, nitrification inhibition). Literature sources used in this study were selected based on the following criteria: (1) The research object was swine; (2) The study included at least one of the CH₄, N₂O and NH₃ gases; (3) Gas emission flux or gas emission factor was available; (4) For literature related to mitigation, only studies that reported at least one control group were selected so that emission mitigation efficiency could be calculated.

Data and analysis

The Wilcoxon Signed-Rank test was used to determine if the median mitigation efficiency was significantly different from zero when there were sufficient results for specific measures.

Number of papers	Population	Intervention	Comparator	Outcome	Quality score
142	Swine manure	Anaerobic digestion; Composting with additives	No mitigation strategy	Metric: NH ₃ emissions; Effect size: Ratio of the considered metrics in the intervention to the considered metrics in the control	62.5

Results

- Land application of digested slurry as compared to raw manure was not efficient in reducing NH₃ emissions ($p > 0.05$).
- Adding additives before composting manure reduced NH₃ emissions by 42 (P=0.028).

Factors influencing effect sizes

- No factors influencing effect sizes to report

Conclusion

Land application of digested slurry as compared to raw manure was not efficient in reducing NH₃ emissions ($p > 0.05$). Composting with additives significantly reduced NH₃ emissions.