

## IMPACT: GHG EMISSIONS

### Reference 8

Wang, Y; Dong, HM; Zhu, ZP; Gerber, PJ; Xin, HW; Smith, P; Opio, C; Steinfeld, H; Chadwick, D 2017 Mitigating Greenhouse Gas and Ammonia Emissions from Swine Manure Management: A System Analysis ENVIRONMENTAL SCIENCE & TECHNOLOGY 10.1021/acs.est.6b06430

### Background and objective

Studies have been conducted to address manure-related emissions, and various mitigation measures have been tested and developed. However, most studies have focused either on one specific gas, one individual manure management phase or influencing factor, or mitigation practice. The objective of this study is to estimate the emissions mitigation potentials for NH<sub>3</sub>, methane (CH<sub>4</sub>), and nitrous oxide (N<sub>2</sub>O) of different swine manure storage and treatment mitigation strategies. Here the results concerning CH<sub>4</sub> and N<sub>2</sub>O emissions are reported.

### Search strategy and selection criteria

The ISI Web of Knowledge database ([www.isiwebofknowledge.com](http://www.isiwebofknowledge.com)) and the Chinese journal database ([www.cnki.net](http://www.cnki.net)) were used to search all published data sets as of January 2016. Specific search terms were combined and used, depending on animal categories (swine, pig, livestock, animal), manure, in-house manure management (slatted floor, pit, bedding, litter, pull-plug, discharge, scraper, separation), outdoor manure management (lagoon, slurry pond, storage tank, compost, solid storage, stockpile), land application (surface spreading, injection, incorporation, band spreading), gaseous emission (NH<sub>3</sub>, CH<sub>4</sub>, N<sub>2</sub>O, and GHG gas), and mitigation measure (diet, biofilter, biogas, additive, cover, acid, cooling, nitrification inhibition). Literature sources used in this study were selected based on the following criteria: (1) The research object was swine; (2) The study included at least one of the CH<sub>4</sub>, N<sub>2</sub>O and NH<sub>3</sub> gases; (3) Gas emission flux or gas emission factor was available; (4) For literature related to mitigation, only studies that reported at least one control group were selected so that emission mitigation efficiency could be calculated.

### Data and analysis

The Wilcoxon Signed-Rank test was used to determine if the median mitigation efficiency was significantly different from zero when there were sufficient results for specific measures.

Number of papers	Population	Intervention	Comparator	Outcome	Quality score
142	Swine manure	Slurry injection; Slurry incorporation; Solid incorporation; Digested slurry; Land application with nitrification inhibitor; Avoiding manure application to rice paddy fields	No mitigation strategy	Metric: 1) CH <sub>4</sub> emissions; 2) N <sub>2</sub> O emissions; Effect size: Ratio of the considered metrics in the intervention to the considered metrics in the control	68.75

### Results

- Avoiding manure application to rice paddy fields is an effective GHG mitigation option, with CH<sub>4</sub> and N<sub>2</sub>O mitigation efficacy of 57% ( $p < 0.001$ ) and 23% ( $p = 0.575$ ), respectively.
- Slurry injection significantly increased N<sub>2</sub>O emissions (+84%,  $p=0.003$ ), while addition of nitrification inhibitors mitigated N<sub>2</sub>O emissions (-28%,  $p=0.016$ ). Other upland application strategies (Slurry incorporation; Solid incorporation; Land application of digested slurry;) were not effective in reducing N<sub>2</sub>O emissions ( $p>0.05$ ).
- NULL
- NULL
- NULL

### Factors influencing effect sizes

- No factors influencing effect sizes to report

### Conclusion

This study shows that avoiding to spread swine manure in rice paddies and adding nitrification inhibitors in the manure before spreading in upland were effective in mitigating CH<sub>4</sub> and N<sub>2</sub>O emissions, while slurry injection increased N<sub>2</sub>O emissions. Land application of digestate, Slurry and solid incorporation showed non-statistically significant effects on N<sub>2</sub>O emissions.