

SINGLE-IMPACT FICHE COVER AND CATCH CROPS

IMPACT: BIODIVERSITY

Data extracted in January 2022 Fiche created in February 2024

Note to the reader: This fiche summarises the effects of Cover and catch crops on BIODIVERSITY. It is based on 1 synthesis paper¹ containing 74 primary studies.

1. WEIGHT OF THE EVIDENCE

CONSISTENCY OF THE IMPACT

The effect of cover/catch crops on BIODIVERSITY is reported in **Table 1**.

The table below shows the number of synthesis papers with statistical tests reporting i) a significant difference between the Intervention and the Comparator, that is to say, a significant statistical effect, which can be positive or negative; or ii) a non-statistically significant difference between the Intervention and the Comparator. In addition, we include, if any, the number of synthesis papers reporting relevant results but without statistical test of the effects. Details on the quality assessment of the synthesis papers can be found in the methodology section of this WIKI.

 The effect of cover/catch crops, as compared to bare soil on BIODIVERSITY is positive, according to 1 synthesis paper assessing plant species richness, earthworm species richness, spider species richness and abundance, beetle species richness and abundance, grasshopper species richness and bird species richness.

The selected synthesis paper included studies conducted in Europe (see Table 2).

Table 1: Summary of effects. Number of synthesis papers reporting positive, negative or non-statistically significant effects on environmental and climate impacts. The number of synthesis papers reporting relevant results but without statistical test of the effects are also provided. When not all the synthesis papers reporting an effect are of high quality, the number of synthesis papers with a quality score of at least 50% is indicated in parentheses. The reference numbers of the synthesis papers reporting each of the effects are provided in **Table 3**.

| | | | | | Non-statistically tested | | |
|-----------------------|--------------|--------------|------------|------------------------|--------------------------|---|---|
| Impact | Metric | Intervention | Comparator | Significantly positive | Non statistically tested | | |
| Increase biodiversity | Biodiversity | Cover crops | Bare soil | 1 | 0 | 0 | 0 |

QUALITY OF THE SYNTHESIS PAPERS

The quality of each synthesis paper was assessed based on 16 criteria regarding three main aspects: 1) the literature search strategy and primary studies selection; 2) the statistical analysis conducted; and 3) the evaluation of potential bias. We assessed whether authors addressed and reported these criteria. Then, a quality score was calculated as the percentage of these 16 criteria properly addressed and reported in each synthesis paper. Details on quality criteria can be found in the methodology section of this WIKI.

2. IMPACTS

The main characteristics and results of the 1 synthesis paper is reported in **Table 2** with the terminology used in those papers, while **Table 3** shows the reference numbers of the synthesis papers reporting for each of the results shown in **Table 1**. Comprehensive information about the results reported in each synthesis paper, in particular about the modulation of effects by factors related to soil, climate and management practices, are provided in the **summaries of the synthesis papers** available in this WIKI.

| Reference number | Population | Scale | Num. papers | Intervention | Comparator | Metric | Conclusion | Quality score |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|----------------|-------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|
| Ref29 | Vineyards. Global dataset. About 40% of all datasets originated from irrigated vineyards, 50% were rainfed vineyards and the other studies did not provide information on the use of irrigation. Most datasets came from vineyards under Mediterranean climates (n = 100), oceanic climates (n = 56), and steppe or continental climates (n = 2; three studies included vineyards from different climates). Most studies implemented randomized block designs within one experimental vineyard (n = 113), only few studies implemented block designs in several vineyards (n = 12), whereas 56 datasets used individual vineyards as replicate. The | Global. Major wine producing regions world- wide except Asian countries, New Zealand and Argentina | 74 | Cover crops or natural vegetation growth for soil cover in vineyards | Bare soil or removal of spontaneous vegetation in vineyards by herbicides use or tillage | Plant species richness, Fauna, Earthworm species richness, Spider species richness and abundance, Beetle species richness and abundance, Grasshopper species richness, Insect pollinator species richness and abundance (bees, butterflies), Bird species richness | Biodiversity significantly increased (53%) and benefitted most from Cover crops or natural vegetation growth for soil cover in vineyards. | 94% |

 Table 2: Main characteristics of the synthesis paper reporting effects on biodiversity.

¹ Synthesis research papers include either meta-analysis or systematic reviews with quantitative results. Details can be found in the methodology section of the WIKI.

| Reference number | Population | Scale | Num. papers | Intervention | Comparator | Metric | Conclusion | Quality score |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|----------------|--------------|------------|--------|------------|------------------|
| | majority of studies investigated the effects of bare soil management (mostly due to tillage, sometimes by use of herbicides or both) compared to cover crops or natural vegetation (n = 137 datasets). We investigated the effects of conventional vs. organic management in 27 studies and 17 datasets originated from other types of intensive vs. extensive vegetation management like the contrast of single to diverse cover crop species in inter- rows or mulching vs. mowing of vegetation. | | | | | | | |

Table 3: Reference numbers of the synthesis papers reporting for each of the results shown in Table 1.

| | _ | _ | - | | Non-statistically tested | | | |
|-----------------------|--------------|--------------|------------|------------------------|--------------------------|-----------------|--|--|
| Impact | Metric | Intervention | Comparator | Significantly positive | Significantly negative | Non-significant | | |
| Increase biodiversity | Biodiversity | Cover crops | Bare soil | Ref29 | | | | |

3. FACTORS INFLUENCING THE EFFECTS ON BIODIVERSITY

No factors were found.

4. KNOWLEDGE GAPS

The authors did not report knowledge gaps in the reviewed synthesis papers.

5. SYNTHESIS PAPERS INCLUDED IN THE REVIEW

Table 6: List of synthesis papers included in this review. More details can be found in the summaries of the meta-analyses.

| Ref Num | Author(s) | Year | Title | Journal | DOI |
|------------|-----------------------------------------------------------------------------------------------------------------------------------------------|------|-----------------------------------------------------------------------------------------------------------------|----------------|-----------------------------|
| Ref29 | Winter, S; Bauer, T; Strauss, P; Kratschmer, S; Paredes, D; Popescu, D; Landa, B; Guzman, G; Gomez, JA; Guernion, M; Zaller, JG; Batary, P | 2018 | Effects of vegetation management intensity on biodiversity and ecosystem services in vineyards: A meta-analysis | J APPL ECOL | 10.1111/1365- 2664.13124 |

2

Disclaimer: These fiches present a large amount of scientific knowledge synthesised to assess farming practices impacts on the environment, climate and productivity. The European Commission maintains this WIKI to enhance public access to information about its initiatives. Our goal is to keep this information timely and accurate. If errors are brought to our attention, we will try to correct them. However, the Commission accepts no responsibility or liability whatsoever with regard to the information on these fiches and WIKI.

3